

Original Research

## Comparative Assessment of Seasonal Variations in the Quality of Surface Water and Its Associated Health Hazards in Gold Mining Areas of Osun State, South-West Nigeria

Awogbami Stephen Olalekan <sup>1</sup>, Solomon Olayinka Adewoye <sup>2</sup>, Sawyerr Olawale Henry <sup>1</sup>, Morufu Olalekan Raimi <sup>3,\*</sup>

1. Department of Environmental Health, Faculty of Pure and Applied Science, Kwara State University, Malete, Kwara State, Nigeria; E-Mails: [awogbamistev@gmail.com](mailto:awogbamistev@gmail.com); [henry.sawyerr@kwasu.edu.ng](mailto:henry.sawyerr@kwasu.edu.ng)
2. Department of Pure and Applied Biology, Ladoke Akintola University of Technology, Ogbomosho, Nigeria; E-Mail: [soadewoye@lautech.edu.ng](mailto:soadewoye@lautech.edu.ng)
3. Department of Community Medicine, Faculty of Clinical Sciences, Niger Delta University, Wilberforce Island, Bayelsa State, Nigeria; E-Mail: [morufuolalekan.raimi@gmail.com](mailto:morufuolalekan.raimi@gmail.com)

\* **Correspondence:** Morufu Olalekan Raimi; E-Mail: [morufuolalekan.raimi@gmail.com](mailto:morufuolalekan.raimi@gmail.com)

**Academic Editor:** Wen-Cheng Liu

**Special Issue:** [Advances in Hydrology, Water Quality and Sediment Simulation Modelling](#)

*Adv Environ Eng Res*

2023, volume 4, issue 1

doi:10.21926/aeer.2301011

**Received:** October 30, 2022

**Accepted:** January 05, 2023

**Published:** January 23, 2023

### Abstract

Industrial and urban development are contributing to an increase in global environmental degradation. Therefore, the release of heavy metals from mining-related operations into surface water is harmful to human health. But as anthropogenic influences on the environment grow, surface water characteristics are also alter, impacting aquatic creatures. In order to analyze the acceptability of the surface water in the gold mining area for drinking and irrigation uses, different quality water assessment methodologies were incorporated in this research. In light of this, the purpose of the current study is to comprehend how the hydrogeochemistry and appropriateness of surface water for drinking and irrigation vary



© 2023 by the author. This is an open access article distributed under the conditions of the [Creative Commons by Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium or format, provided the original work is correctly cited.

every month. The study employed standardized analytical techniques. According to APHA recommendations, all sampling, conservation, transportation, and analysis were completed (2012). All collected samples were transported to the study lab while being kept in an icebox to prevent the degradation of the organic components. As a result, the study is focused on the contamination level in the surface water for a year. Overall, the study highlights important pollutants impacting surface water quality as it passes through Osun State's gold mining regions. Finally, it has been determined that the following criteria are crucial for the stretch of this research season: DO, Hardness, Turbidity, Chloride, Potassium, Nitrate, Lead, TSS, Cadmium, Chromium, Sulphate, Manganese, Mercury, and Arsenic. Most of the physicochemical variables examined in this study fell within their corresponding standard limits. Based on the results of this study, the appropriate constituted authority is encouraged to continuously monitor and assess surface water quality suitability for drinking, domestic, and irrigation purposes by keeping track of the effects of water contaminants and detecting any changes in the water quality. To safeguard and maintain groundwater quality and public health, it is advised that appropriate regulatory policies and water treatment procedures be employed in the area. Additionally, it is proposed that when enhancing water quality and investigating the sustainable use of water resources, surface water pollution should be taken into consideration.

### **Keywords**

Hydrochemistry; sustainability assessment; drinking water quality; physicochemical; irrigation water quality; artisanal and small-scale gold mining (ASGM); emerging contaminants; South-West Nigeria

## **1. Introduction**

Nigeria is fortunate to have a wealth of water resources, with an estimated 226 billion m<sup>3</sup> of surface water and 40 billion m<sup>3</sup> or so of groundwater. According to WHO guidelines, 28% of the world's population lives in Africa and doesn't have access to better water supplies. Water quality is the main factor limiting aquatic ecosystems' productivity, particularly fish resources. According to the United Nations Educational, Social, and Cultural Organization, water supply is essential in both rural and urban settings. Water is a common element to the other four fundamental human needs (food, health, education, and peace), making it essential to life and human growth. Water is crucial in developing settlements and determining population density [1-10]. Furthermore, surface waters, like shallow lakes, are dynamic systems with a significant spatiotemporal variation. Lithology, water flow rate, geochemical reaction types, salt solubility, and human activities often control the differences in the concentration of dissolved ions in surface water. Eutrophication, which degrades water quality by encouraging excessive algal growth and elevating the concentration of suspended organic material and heavy metals, is currently one of the most prevalent ecological issues affecting inland water bodies [11-16]. Even though there is a lot of water on the earth, usually surface and groundwater and it is distributed through a variety of media, including rivers, lakes, oceans, seas, glaciers, and streams [1-21], only around 0.3% of it is

potable and used by people [22-25]. The fact that water doesn't exist in isolation and always has some level of contamination due to its property as a universal solvent depends on several variables, including the geology of the environment or the nature of aquifers, anthropogenic activities occurring nearby, and the level of sanitation and hygienic standards in the community in question [14-16, 25-27]. The increased demand for potable water for domestic, industrial, agricultural, and recreational uses, as well as other anthropogenic activities, has been linked to rapid industrialization, urbanization, and population growth worldwide [28-39]. This has resulted in a significant burden of water contamination. Surface water makes up the remaining percentage at the same time. The quantity and quality of water suitable for drinking, residential use, and commercial use are declining. The assessment and monitoring of water quality for various reasons to conserve, sustain, and protect water resources has piqued the interest of scholars in numerous disciplines of study across the globe [1, 8-13, 18]. Surface water contamination, however, is a major focus of study and research because of the extensive industrialization and urbanization. Water contamination risks have seriously caused the rivers to decline, disrupting the aquatic ecosystem, especially in developing nations like Nigeria. Water contamination is caused by small-scale and artisanal gold mining. This is not implausible given that gold mining operations, particularly alluvial gold mining, take place on the river bed [40]. Waste products from gold mining or processing are frequently discarded or washed into local water sources by rain or flooding. These contaminants significantly impact water quality's physical, chemical, and biological indices. When waste from gold mining activities is dumped into water bodies, it causes an increase in both turbidity and total suspended solids. This has led to the widespread adoption of these measures as markers of water pollution in mining sites [41]. As a result, as the water's temperature increased, the amount of dissolved oxygen decreased [41]. In other places, total dissolved solids and turbidity were the main indicators of artisanal mining contamination [42]. A notorious source of water pollution is mining. One of the main causes of the world's declining water supplies is mining poisoning of water bodies [43]. The unorganized informal mining sector known as artisanal mining has today increased the environmental effects of mining [44]. The minerals are extracted using localized, affordable, and simple-to-use instruments and equipment in artisanal mining [45]. Most of this country's poorest regions engage in this illegal mining sector, which typically operates outside its legal and regulatory framework [46, 47]. Rapid population expansion, industrialisation, and urbanization all worsen this scenario by increasing demand for and pressure on mineral resources [48]. The delicate ecological systems' equilibrium, including the biotic and abiotic components of the environment, are altered by mining operations [49]. Land, soils, and water are contaminated and degraded due to gold mining [28, 29, 50-57]. In addition, heavy metals are present in the wastewater from ore processing and other mining effluents, contaminating water sources [58, 59]. High levels of heavy metals in rivers and streams are likely to bioaccumulate in fish and other aquatic life, posing a major health risk to humans who eat those things [60, 61]. As a result, many rural communities in Nigeria struggle to get access to drinkable water [1-21, 62, 63]. This led to a dependence on freshwater resources, such as rivers and streams, for residential water needs [25]. This claim was supported by a wide range of reliable scientific discoveries and evidence, all of which revolved around the need for water as a necessary component of survival [1-25]. Water is essential for human life and is important for a country's economic prosperity, well-being, and overall health [64-66]. In the meantime, heavy metal contamination is well documented in places where gold is mined. The bioaccumulation of these heavy metals in the

human body can result in diseases such as kidney damage, cancer, reproductive disorders, brain damage, spontaneous abortion, etc. [1-25]. Therefore, it is crucial to research the heavy metal and Physico-chemical composition of water sources in gold mining towns to safeguard public health from potential chemical exposure risks. Additionally, this will give decision-makers in both governmental and non-governmental groups the crucial data they need to take action. As a result, the region considered for this research project is dominated by urban centers, mining industries, and agricultural lands. This work has been attempted with the main objectives of evaluating surface water aptness for irrigation and drinking practices, by computing various monthly variations in Physico-chemical, radical, and heavy metal concentration in selected surface water bodies. As there is no known extensive research study carried out so far in this study area for assessing surface water quality and its associated health hazard in a gold mining area of Osun State, this work has been attempted. The results of this study may help the locals use surface water resources wisely for beneficial agricultural practices. In agricultural settlements in Nigeria, the quality of the surface water has been in issue. The majority of these surface waters, according to reports, are unsafe for drinking and irrigation. However, some may be suitable for industrial uses. According to Raimi and Sawyerr [11], background values for some components in the oil and gas environment are assessed either spatially or temporally (concentrations prior to anthropogenic activity) (concentrations in the areas not influenced by anthropogenic activity). As a result, there is an increasing need to address surface water quality issues because poor water quality impacts both crop output and human health. Anthropogenic and geogenic factors significantly impact changes in surface water quality [7, 9, 11, 12, 17]. Therefore, if anthropogenic activities are to blame for the degradation, environmental legislation can preserve surface water. Monitoring the river's water quality is also crucial for sustainable use. However, long-term monitoring produces a huge and complex database that requires a competent method for interpretation. To better understand geographical changes in water quality, multidimensional scaling analysis will aid in interpreting of complicated datasets. These methods are excellent resources for creating sensible plans for the efficient management of water resources. Hence, this paper presents a new set of data on surface water geochemistry from gold mining areas of Osun State, South-West Nigeria.

## **2. Material and Methods**

Surface water samples analyzed in this research were collected from three surface water bodies from different locations for twelve (12) months (June 2020 to May 2021). The surface water bodies include the Aye-Oba River, Alapadi and Eti-oni stream; both Alapadi and Eti-oni stream are tributaries of the Aye-oba River. All are situated within gold mining activities areas across three Local Governments (Ife South, Atakumosa East, and Atakumosa West) in Osun State, Southwest Nigeria. Aye-Oba River lies between the latitude of 7°23' N and the longitude of 4°61' E (see Figures 1-5). The river is very significant to the socio-economic growth of the residents of Ife South Local Government area in Osun State to sustain subsistence farming such as irrigation, livestock, and fishery. The river is the major source of water supply for consumption and domestic use. The river was constructed as a dam in the year 2004. Thus, the dam served as the major source of potable water supply to more than fifteen communities in Ife South Local Government areas and its environs until it stopped operation (Figures 1-5).




Figure 1 Area map showing sampling Aye-Oba River and its tributaries.




Figure 2 Map showing sampling points at Aye-oba River and in Ife South.




Figure 3 Map showing sampling points and LULC Classes in Ife South.




Figure 4 Map showing sampling sites in Atakumosa East.




Figure 5 Map showing sampling points in Atakumosa West.

### 2.1 Experimental Design and Description of the Sampling Points


The study design employed was a laboratory-based experimental approach and a descriptive cross-sectional study coupled with field observations. The sampling points were: upper stream (site A), middle stream (site B) and lower stream (site C) (see Figures 1-5 above). The division was

done concerning the discharge of effluents from mining, domestic and agricultural activities, especially pesticide pollution through various forms of indiscriminate application on farmlands and domestic effluents particularly from the oil mills processing that enter the stream. Site A (upper stream) represents the Eti-oni stream; this stream is about 500 meters from the mining site and about 1 km to site B the effluent discharge point.

Site B (middle stream) represents the Alapadi stream. This stream directly receives effluent discharge due to mining activities in the area. It also receives significant effluents from oil palm processing units along the river bank and runoff water from farmlands, cassava processes, and refuse dumps from homes. Site C (lower stream) represents the Aye-Oba River/dam, the abandoned dam. The river is about three (3) kilometers from site B. Physical observation showed that the effluents received by the water bodies from sites A and B changed the color of the water bodies and there was an unpleasant smell emanating from the river. In addition, the river receives effluents from cassava processing wastes, open defecation, poultry wastes, animal dung, pesticide products, and water run-off from the dump site. Therefore, the choice of the aforementioned sampling points, presented in Figure 1-5, was based on the accessibility, the rate at which they receive effluents from different sources, the extent of their pollution, and particularly their distances from the site of mining activities. In addition, sampling sites were chosen based on the potential exposure of the surface water to different agro-industrial and other sources of pollutants.

## ***2.2 Water Sample Collection for Heavy Metal Analysis***

Water samples were collected between 8.00 am and 12 noon, every fourth week of the month in all the sampling points for 12 months (June 2020 to May 2021). Water samples were collected into plastic bottles which were previously soaked in 3% nitric acid and washed with distilled water before sampling, this was in accordance to the method described by APHA [67]. Water samples for the determination of dissolved oxygen were collected in dark glass containers and fixed on the spot with a Winkler reagent. The water samples were properly preserved following the water sample preservation methods described by APHA [67]. As a result, Figure 6 displays the primary approaches for determining the composition of surface water.



**Figure 6** A diagram depicting the quantification methods used in the current investigation. Adapted from Raimi *et al.*, [9], Olalekan *et al.*, [10], Raimi & Sawyerr [11], Raimi *et al.*, [12].

### 3. Results and Discussion


#### 3.1 General Description of Monthly Variations of Surface Water Quality

Understanding a water body's health depends critically on its water quality. Numerous researchers have shown over the past few decades that surface water interfaces' physical and chemical processes are extremely complicated and frequently interconnected. Surface water pollution by heavy metals is a significant environmental problem, and numerous studies have been conducted to identify rising metal concentrations that cause increased toxicity. This leads to a sharp decline in microbial activity, which is reflected in a slowing of the apparent growth rate and an extension of the lag time. Figures 7 through 34 below show a descriptive summary of all the analyzed physicochemical parameters. Poor mining practices have been used in the study area, primarily agricultural. It is thought that several pollutants may have impacted this region's surface water resource. The next section includes several physical chemicals that are considered to be "good water." While the study documents the physicochemical characteristics of the study area's surface water. The significance of the study is to evaluate the water's appropriateness and quality for home use. Due to the study area's proximity to a residential area, the river and its tributaries serve as a water source for domestic use. The water body passes via a mine, an abandoned dam, a processing plant for oil palm, a processing plant for cassava, and farms with cattle and fisheries.

Potable water must be flavorless, odorless, and colorless, by Morufu and Clinton [1], Olalekan *et al.* [10], and other sources. These specifications weren't met by the surface water used in this


study. These might occur from microbial activity in the water. According to reports, biological processes and chemical contamination of water sources encourage the development of microbial communities and give water an unpleasant odor, look, and taste [13]. Due to its role in bodily processes, water is the most crucial nutrient for human survival, according to Raimi *et al.* [2].

Similarly, the author claimed that water is crucial to human nutrition, directly as drinking water and indirectly as a food medium in addition to its many other uses. Lack of access to adequate drinking water has been linked to a number of health issues that plague developing countries like Nigeria [1-25]. About 66.3 million Nigerians [13-16, 19, 24-27] lack access to clean drinking water, which causes them to rely on surface water for their daily needs and as a method of waste disposal. According to studies by Olalekan *et al.* [14], Raimi *et al.* [16], Olalekan *et al.* [17], and Raimi *et al.* [19], there are about 2 billion people without access to potable drinking water in their homes worldwide as of 2017. Nearly 80% of these people rely on surface water, which is unsafe for drinking as well as other domestic uses. Olalekan *et al.* [10] assert that pure water is colorless. Therefore, any water with a distinctive color implies pollution. Figure 7 shows the monthly fluctuation in surface water's apparent color. In January (1925.35), June (2030.18), July (1920.28), September (1720.52), and November, the apparent color of the surface water was high and over 1500. (1568.57). In surface water, March saw the lowest value of 447.07 and June saw the highest value of 2030.18. This study is in opposition to the study that discovered that Okpai had the highest value for color during the dry season, with a value of 35.33, while Okpai had the lowest value during the wet season, with a value of 32. The rainy season must have had a greater impact on the color of surface waters than the dry season. Surface water is typically highly contaminated during the rainy season.



**Figure 7** Monthly variation of apparent colour in surface water.


The receiving water bodies may experience significant DO depletion and fish deaths due to a high biochemical oxygen demand (BOD) level [12]. The BOD measures the quantity of organic matter in biologically active water. Figure 8 displays the monthly variance in the surface water's BOD. January, April, and the early to late rainy seasons saw elevated BOD levels in surface water (June–September). The maximum BOD in surface water was recorded in August (4.4 mg/L), while the lowest value was recorded at the start of the rainy season in May (0.93 mg/L). High BOD levels were also recorded in January (4.27 mg/L) and June (4.00 mg/L). BOD<sub>5</sub> is a crucial measure for detecting contamination from organic and inorganic wastes. As a result, wet seasons had higher values than dry ones, leading to the conclusion that anthropogenic activities may have an impact on higher BOD levels in the same way wetness in seasonality had a greater impact on BOD than dry seasons. Therefore, the discharge into surface water from industrial units engaged in gold mining accounts for this rise in BOD<sub>5</sub> concentrations. If no action is taken to address BOD<sub>5</sub> trends, the environment of the area receiving these effluents will suffer grave effects. BOD<sub>5</sub> must not exceed the 40 mg/l limits for discharge into the environment. But since the BOD<sub>5</sub> concentrations in the surface water have gone above the permitted environmental release threshold, careful action is needed to lessen the effects of this pollution.



**Figure 8** Monthly variation of Biological Oxygen Demand (BOD) in Surface water.

COD are significant indicators of organic and inorganic waste pollution [1-6]. Higher COD values were linked by Raimi *et al.* [12] to greater anthropogenic stresses on groundwater. As a result, Figure 9 provides the chemical oxygen requirement in surface water. For instance, the mid-late dry season (January–April) and the middle of the rainy season have greater chemical oxygen demand levels (June–August). The highest value (9.60 mg/L) and lowest (3.20 mg/L) were noted in April and July, respectively. Chemical oxygen demand levels averaged 7.47, 4.27, and 5.60 mg/L in January, February, and March, respectively, while they were 6.93, 6.67, 5.07, 5.07, 6.67, 6.67, and

6.13 mg/L in May, June, August, September, October, November, and December. Thus, it might be concluded that locations susceptible to mining operations have a greater influence on COD than those not affected by these activities. Additionally, it might be proven that rainy seasons had a greater impact on COD than their dry counterparts. Higher COD values were linked by Olalekan *et al.* [10] and Raimi *et al.* [9] to greater anthropogenic stresses on groundwater. However, urgent action must be taken to lessen the impact of this flow's environmental pollution. 150 mg/l is the upper limit for COD discharge into the environment. The COD trends in the surface water channels, on the other hand, are obviously below 150 mg/l.




**Figure 9** Monthly variation of Chemical Oxygen Demand (COD) in Surface water.

Aquatic creatures require dissolved oxygen (DO) for their best chances of survival. Low oxygen levels are a sign of biological activity, nutritional input, and organic loading [6, 13]. Large loads of organic debris are frequently the source of elevated water DO values. In extreme circumstances, oxygen loss can result in major fish death by altering the fish population [25, 31]. For good fish production, a DO level of at least 5 mg/L is advised [6, 13, 31]. So deviation from that range impacts fish survival in that body of water. Its concentration in surface water fluctuates depending on the trophic levels of the water. The amount of dissolved oxygen is influenced by photosynthetic activity and the microbial breakdown of both native and foreign organic materials. The surface water's generally low level of dissolved oxygen indicates eutrophication. The most frequent outcome of several types of water pollution is likely the depletion of DO in the water [17, 18, 20, 68]. As a result, Figure 10 provides the DO concentrations in surface water. Surface water DO changes by month show some obvious differences.

Compared to the dry season (5.33–7.6 mg/L), the DO levels in surface water are greater during the rainy season (6.4–9.6 mg/L). The months with the greatest DO levels were June (9.6 mg/L) and September (9.07 mg/L), while the months with the lowest levels were March (5.33 mg/L). As a result, higher DO correspond to increased biological activity; rainy seasons had a greater impact than dry seasons. However, there were no discernible variations across the different months at the p0.05 level of significance. The pH and dissolved oxygen (DO) levels, however, were within the WHO-recommended range [69] and matched the findings of Afolabi and Raimi [21]. The suggested

value was not exceeded by any of the chemical ions extracted from this study's water samples. According to the study, they are less than the WHO offers [69]. DO is therefore crucial for supporting a variety of aquatic life. Its concentration in rivers fluctuates depending on the trophic levels of the lakes. The amount of dissolved oxygen is influenced by photosynthetic activity and the microbial breakdown of both native and foreign organic materials. The river's overall low level of dissolved oxygen points to eutrophication. Certain types of water pollution often lead to the depletion of DO in the water.


Additionally, the current trends of DO depletion in the majority of sample stations are brought on by the existence of a large organic load, dumped by a drain, and religious rituals along the river bank. Hydrogen sulfide, ammonia, nitrite, ferrous iron, and several oxidizable compounds are inorganic reducing agents that lessen the amount of dissolved oxygen in the water. Meanwhile, decomposing extra nutrients and biodegradable organic materials by decomposing organisms like bacteria may cause the low DO value during the dry season. These organic materials are brought in by an influx of dissolved solutes from nearby metropolitan areas, agricultural fields, and industrial wastes.



**Figure 10** Monthly variation of Dissolved Oxygen (DO) in surface water.


Figure 11 displays the electrical conductivity (EC) level in surface water. Surface water EC values were typically higher from February (206.67 S/cm) and March (210.00 S/cm) through April (224.90 S/cm) and May (224.97 S/cm) until the end of the dry season and the beginning of the rainy season. As a result, it considers the nutrient load of rivers, which are highly impacted by anthropogenic activities such as wastewater discharges and agricultural runoff. May saw the highest EC value of 224.97 S/cm, while July saw the lowest value at 103.33 S/cm. As a result, an increase in EC in water can result in aesthetic issues and annoyances, such as an unwelcome taste and color [22]. The 500–600 S/cm range corresponds to the WHO's recommended threshold for drinking water [22]. Afolabi and Raimi [13] asserted that poor water quality is not related to a greater level of EC. In addition, the presence of dissolved salts and other organic resources may be the cause of increased conductivity readings. Olalekan *et al.* [25] also noted that conductivity readings higher than 100 S/cm showed human activity. Water conductivity between 150 and 500

S/cm is optimum for fish culture, according to Olalekan *et al.* [10]. This showed that the EC values found in the current study were higher than those considered ideal for fish culture. This could be due to the high conductivity agricultural drainage, the solutions of most inorganic compounds, and the more numerous ions produced by industry [67]. This could be due to the high conductivity agricultural drainage, the solutions of most inorganic compounds, and the more numerous ions produced by industry [67].




**Figure 11** Monthly variation of electrical conductivity (EC) in Surface water.

Figure 12 depicts the surface water's hardness. Surface water's hardness peaked in April at 182.94 mg/L before dropping precipitously and reaching its lowest point in June (7.54 mg/L). The readings for July (73.21 mg/L), August (71.61 mg/L), September (89.30 mg/L), October (74.55 mg/L), November (74.90 mg/L), and December (78.13 mg/L) showed little variation. Alkaline earth, which includes calcium and magnesium ions, causes water's total hardness (TH) (it measures the sum of calcium and magnesium ions). According to the classification system, water with a TH of less than 60 mg/L is categorized as soft, 60 to 120 mg/L is moderately hard, 120 to 180 mg/L is hard, and more than 180 mg/L is very hard (Figure 12). The water used for this investigation is categorized as soft and moderately hard according to the chart. According to this study's TH values, divalent metallic ions, calcium, and magnesium ions are dissolved in low to moderate amounts [9-12].




**Figure 12** Monthly variation of hardness in Surface water.

Total dissolved solids (TDS) originate from natural sources, sewage, urban runoff, and industrial wastewater (WHO, 2017). A high level of TDS affects aquatic life (APHA, 2012). Thus, higher ionic concentration, which is less palatable and causes an undesirable physicochemical reaction in consumers, is indicated by high TDS in water [6, 13]. TDS levels in surface water were measured between 89 mg/l and 158 mg/l (Figure 13), with the lowest value occurring in the rainy (monsoon) month of July (89 mg/l) and the highest in the dry season of March (158 mg/l). Due to the significant concentration of dissolved organic matter and dissociated electrolytes entering the surface water through various point and non-point sources, upward trends in TDS were seen during each monitoring month.




**Figure 13** Monthly variation of total dissolved solids (TDS) in surface water.

The total organic carbon (TOC) and total organic matter (TOM) level in the surface water is given in Figure 14. The TOC and TOM trends were identical as TOM was estimated from TOC. The peak values of TOC were recorded between June (21.88 mg/L) and July (21.04 mg/L). In the mid-late dry season, the range of TOC was between 3.04–7.11 mg/L. In the rainy season, it was between 6.58–21.88 mg/L. For TOM, highest value was observed in June (38.0 mg/L), July (36.5 mg/L), April (10.0 mg/L) during rainy season. While in January, October, November and December the values are 10.2, 10.2, 10.2 and 10.0 mg/L respectively.



**Figure 14** Monthly variation of total organic carbon (TOC) and total organic matter (TOM) of surface water.


The true color of surface water is given in Figure 15. The values were higher for most of the study period, particularly mid-dry season (January) and Mid-rainy season (July), where it peaked at 1037.73 TCU and 1331.28 TCU respectively. Its lowest value was in March (253.01 TCU). The value of surface water's true color decreases toward the beginning of the dry season in October (294.37 TCU). Several chemical, physical, radiological and biological parameters have characterized good and potable drinking water. In general, drinking water's appearance, taste, color, and odor are used to determine its quality [9-12]. Hence, the true colour of water samples in the present study were extremely high in January, June, July and September and when compared to the WHO standard; this may be due to effluents from industries, mining activities and homes around the study area [1, 25].



**Figure 15** Monthly variation of true colour in surface water.

Turbidity restricts light penetration and limits photosynthesis in the aquatic environment. The turbidity degree of the water is an approximate measure of the intensity of the pollution [22, 23]. High turbidity indicates the presence of organic suspended material, which promotes the growth of microorganisms [9, 11]. Also, the water turbidity level describes the cloudiness of the water as a result of the precipitation of chemical, suspended particles, faunas and flora debris in the water bodies [29]. Turbidity in the surface water is shown in Figure 16. The turbidity level in surface water in some of the months was markedly higher, most especially in the months of the rainy season, this may be related to flood water originating from surrounding the research area. The highest turbidity value could be the presence of high biodegradable organic matter that comes from wastes of surrounding urban and is discharged from mining-related activities and agricultural fields.


There is an inclination of turbidity levels from June, July, August, and September. Surface water was high in most of the rainy season between, June and September (163.38–222.94 NTU). Its lowest point was recorded in May (35.43 NTU). Thus, turbidity could be due to continuous and impactful predisposition to receiving large quantities of organic and inorganic materials emanating from mining-related activities contaminating the surface waters of the study area. Raimi *et al.*, [9], Olalekan *et al.*, [10], Raimi and Sawyerr [11] and Raimi *et al.*, [12] attributed high values of turbidities in the dry season to decreased vegetation and evapotranspiration during cooler months. Thus, the present study reports high turbidity in all the water samples, making the water unsuitable for human consumption. All the water samples collected along the course of the river, in both dry and wet seasons were higher in surface water, especially in the dry season. This indicates the possibility of the water bodies containing hazardous chemicals and microorganisms (bacteria and protozoa) which are pathogenic to humans [23]. The results of the turbidity level recorded from this study fall within the turbidity value reported by Olalekan *et al.* [25], which is higher than the WHO-recommended value.



**Figure 16** Monthly variation of turbidity in surface water.


Temperature plays a vital role in determining the effectiveness of the fish’s digestive enzymes, reproductive activities, and life cycles [23, 39]. A study by Afolabi and Raimi [13] showed that temperature influences fish growth, specifically in the sensitive fingerling stage. Olalekan *et al.* [25] found that high water temperature is an optimum condition for various mesophilic bacteria to grow, thus playing an important role in influencing their presence in fish. Thus, the water temperatures fluctuate naturally daily and seasonally with air temperature. Surface water bodies can buff water temperature; even moderate changes in water temperature can seriously impact the river ecosystem due to narrow temperature tolerance by aquatic organisms. High amounts of sewage discharges as well as religious ritual activities along the river bank significantly change the river water temperature. Thus, temperature plays a vital role in controlling the chemical and biological composition of a freshwater body. In the aquatic environment, the temperature is the most significant ecological factor. In Osun state, rivers show seasonal temperature variation. The temperature of surface water is given in Figure 17. There was a slight difference in temperature across the months in the surface water. The range of value was 24.93–30.13°C. No clear peak was observed. The slightly low temperature from this study was recorded in July at 24.93°C which is not too obvious from other values of temperature recorded from other months.

Afolabi & Raimi [13] and Odipe *et al.*, [22] stated that areas prone to discharge industrial wastes usually have temperatures above those of their surrounding environments. Thus, the operational presence must have influenced an increase in surface water temperature, correspondingly reflecting in the result as seen above. This indicates surface water pollution since organisms that initially depend on surface water could find the temperature ranges no longer suitable for their continued stay and could migrate to areas with favorable temperature ranges. Moreover, the present study, was higher than the values reported in the studies by Raimi *et al.* [12] and Olalekan *et al.* [14], who reported the  $23.5 \pm 1.8^\circ\text{C}$  and  $21.23^\circ\text{C}$ , respectively. Morufu and Clinton [1] recommended a desirable temperature range of 20-30°C for aquaculture water quality. Afolabi and Morufu [6] also recommended a temperature range of 20 to 35°C for surface water. This indicates that the temperature values were within the recommended limits and that the same temperature range is also sufficient for the proliferation of most pathogenic bacteria [10].




**Figure 17** Monthly variation of surface temperature.

The TSS concentrations in the surface water are given in Figure 18. The levels of TSS were generally low in February (29.17 mg/L), March (80.00 mg/L), and May (24.33 mg/L). However, in the rainy season, they reached a peak of 2547.33 mg/L in September. The range of values was 24.33–2547.33 mg/L. Thus, the excessive influx of suspended solids in surface water could be attributed to the discharge of large quantities of substances directly into surface water bodies or out rightly onto terrestrial areas from where they leach into surface water bodies. Hence, the value of total suspended solids (TSS) reported by Raimi *et al.* [2] is comparable to those of this present study as they both exceeded the recommendation of the WHO guideline for drinking water quality [69].




**Figure 18** Monthly variation of total suspended solid (TSS) in surface water.

The levels of TS in surface water are given in Figure 19. The concentrations in surface water sources are low between the mid-dry season (February)–the early rainy season (June). However, the values increased and peaked in September (2647.33 mg/L), leveling off in the late rainy season to the early dry season. Thus, their research shows that the concentration of solids dissolved in the water determines the concentration of water conductivity. The recommended EC for drinking water according to WHO [69] should not exceed 400  $\mu\text{S}/\text{cm}$ . The EC recorded from this study is found below this value, which agrees with the result published by Odipe *et al.* [22].



**Figure 19** Monthly variation of TS in surface water.


The pH is a general measure of the acidity or alkalinity of a water sample and is indicated on a scale of 0-14. It influences many biological and chemical processes in water. The natural or human-induced process may elevate or decrease the pH of water. Due to its influence on nutrients' solubility and availability as well as their utilization by aquatic organisms, pH becomes an important factor. It varied significantly throughout the seasons. The monthly variations of pH levels in the surface water source are presented in Figure 20. pH level is almost similar through the dry season with a range of 6.10–6.82. However, pH levels of surface water levels increased, reaching a peak of 7.8 in September. Using the maximum permissible range of 6.0-8.5 as the limit for pH as a benchmark [67]. It is seen that the water is acidic. Thus, signifying some level of pollution throughout the seasons. The pH range from 6.10 to 7.8 indicates the productive nature of the water body. This agrees with the discovery by Nwankwo and Ogagarue [70], that areas prone to mining area have pH levels within acidic ranges. In addition, the month of September showed higher acidities during rainy seasons; this could be attributed to large amounts of water received by rainwater which tends to increase the level of acidity within the study area. Hence, it is necessary to take appropriate measures to stop this increase. These results agreed with Olalekan *et al.* [10], Raimi and Sawyerr [11], Raimi *et al.* [12] and Afolabi and Raimi [13] who indicated that pH value lies on the acidic side. Morufu and Clinton [1] concluded that the suitable pH range for aquatic organisms especially for groundwater could be set at 5.5-9.0, implying that the pH value of water recorded during this research was not within the limit, especially during the rainy and dry seasons.



**Figure 20** Monthly variation of hydrogen ion concentration (pH) in surface water.


### 3.2 Monthly Variation of Free Radicals in Surface Water

The  $\text{Ca}^{2+}$  content of the surface water is presented in Figure 21. The range of Calcium ion ( $\text{Ca}^{2+}$ ) in surface water was found between 1.29 mg/L obtained in June and 37.08 mg/L in May. The concentration of  $\text{Ca}^{2+}$  increases from the onset of the dry season toward the onset of the rainy season. There is a decrease in  $\text{Ca}^{2+}$  concentration at the peak of the rainy season. Thus, the presence of Ca and Mg ions in the water supplies is attributed to the occurrence of calcic and ferromagnesian mineral-bearing rocks [70].



**Figure 21** Monthly variation of Ca<sup>2+</sup> in surface water.


Cl<sup>-</sup> levels in the surface water are given in Figure 22. The values of Cl<sup>-</sup> in surface water in the dry season are low. However, surface water Cl<sup>-</sup> increased dramatically in the mid-rainy season (July) reaching a peak of 496.3 mg/L only to plummet in August (21.27 mg/L) and remain steady for the rest of the season. Thus, higher chloride value during the rainy season could be due to large quantities being leached into surface water from adjoining lands due to contaminated rain falling onto such surface water than dry counterparts and settling on such surface waters. Higher chloride values could be due to chloride existing as a natural resource where there was a limited quantity of water to neutralize available chloride compared to the lower value during other months where there are enormous quantities of water to cause massive chloride neutralization. Thus, seasonal variations were also needed wherein it was found that chloride values were higher during the rainy season than during the dry season.



**Figure 22** Monthly variation of Cl<sup>-</sup> in surface water.

Mg<sup>2+</sup> concentrations in surface water are given in Figure 23. In surface water, the highest levels of Mg<sup>2+</sup> were observed between the late dry season and the start of the rainy season, i.e., April (23.80 mg/L) and May (19.07 mg/L). However, the lowest point Mg<sup>2+</sup> concentration was in June


(1.049 mg/L). Thus, it could be deduced that seasonal variations significantly influence magnesium concentrations during both seasons and this seasonal influence was stronger.



**Figure 23** Monthly variation of Mg<sup>2+</sup> in surface water.


NO<sub>3</sub><sup>2-</sup> levels in surface water are shown in Figure 24. NO<sub>3</sub><sup>2-</sup> was low in surface water in both the early rainy season and some months in the dry seasons. Its highest point was toward the end of the rainy season in September (7.37 mg/L) and October (7.82 mg/L). On the other hand, it was somewhat steady between January and August. Nitrates are naturally occurring or anthropogenically incepted environmental pollutants. It is essential for human health but excessive intake may cause adverse health challenges [9-12]. Nitrate is essential in the production of inorganic fertilizers. Its release into water bodies may be through agricultural activities [31, 33-38, 71-73], fossil fuel combustion, and the release of domestic and industrial sewages [4, 7, 8, 12, 74-77]. Methemoglobinemia (also known as a blue baby syndrome) and stomach cancer are associated health hazards of excessive nitrate intake. Also, the high NO<sub>3</sub><sup>-</sup> value could be due to the deamination of ammonium nitrogen from nitrogenous materials and raw wastes that can be oxidized to nitrate by the action of microbiological agents, wastewater disposal, and agricultural activity [2-7, 70]. Henry *et al.* [23] reported that the results of the high rate of microbial activity are associated with a high organic compound and in turn high nitrogen content.

Regarding the toxic nature of NO<sub>3</sub><sup>-</sup>, the World Health Organization (WHO) and the Standard Organization of Nigeria (SON) defined its acceptable limit in water as 50 mg/L. Additionally, seasonal usage of nitrate fertilizers could also explain this trend. The availability of nitrogen-fixing bacteria that penetrate atmospheric nitrogen into the soil could account for the very level of nitrate within the study area and the consequent higher amount in ground waters.




**Figure 24** Monthly variation of NO<sub>3</sub><sup>2-</sup> in surface water.

PO<sub>4</sub><sup>-</sup> is an essential plant nutrient that stimulates the growth of algae and macrophytes in lakes. It is a proxy indicator of lake productivity. PO<sub>4</sub><sup>-</sup> concentrations of the surface water are given in Figure 25. There is fluctuation in the concentrations of the ions throughout the months of the study. However, in July, October and November there was a notable difference between surface water (0.34 mg/L, 0.33 mg/L and 0.34 mg/L were observed). Thus, phosphate groups have been discovered to play a crucial role in the binding of Ni to the cell wall of gram-negative bacteria. PO<sub>4</sub><sup>-</sup> enters the river through domestic wastewater activities accounting for accelerated eutrophication. In addition, phosphate levels were observed to increase from the dry season to the wet season. The rainy season tends to influence phosphate concentration more than the dry season. The higher values in the rainy season at the expense of the dry season could be because farmers in the study area usually engaged in seasonal farming where rain is seasonally targeted before crops could be planted and the soil had to be nourished with fertilizers which phosphate fertilizer is one [31].




**Figure 25** Monthly variation of PO<sub>4</sub><sup>3-</sup> in surface water.

SO<sub>4</sub><sup>2-</sup> levels in surface water are presented in Figure 26. SO<sub>4</sub><sup>2-</sup> concentrations in surface water reached their peak in June (12.50 mg/L) and lowest point in December (2.36 mg/L). It was low at the onset of the rainy season in April (2.87 mg/L) and toward the mid of the dry season in December (2.36 mg/L) and January (2.56 mg/L). Thus, it could be stated that sulfate is very unstable in the atmosphere from where they are converted into forms suitable for its stay in surface and groundwater. Additionally, it could be stated that agricultural contamination from fertilizers seeped underground to mix with groundwater, and gold mining in the study area must have increased the concentration of sulfate during the rainy season as against the low levels during the dry season. Hence, surface water receives ends from rain constituents and contaminants from gold mining activities. This shows the relationship between rain and surface water within the study area.



**Figure 26** Monthly variation of SO<sub>4</sub><sup>2-</sup> in surface water.


Na<sup>+</sup> levels in surface water are given in Figure 27. The surface water had higher Na<sup>+</sup> in the dry season, where it reached a peak of 18.2 mg/L in December and January, its lowest point in July (3.8 mg/L). The range of values of surface water was 3.3–18.17 mg/L and no clear peak was observed. Thus, as seen in the graphs below (Figure 27), the concentrations is low between July, August, October and November but start to rise sharply until December–June, thence to plummet, thus giving a more or less symmetric shape. The reason behind this is the adequate water flow in the river during the rainy season. Quick decisions should be taken to reduce this concentration for environmental protection and public health.



**Figure 27** Monthly variation of Na<sup>+</sup> in surface water.

In Figure 28, the K<sup>+</sup> concentrations in surface water are shown. In both the dry and wet seasons, the K<sup>+</sup> concentrations in surface water are less than 10 mg/L, except the dry season's February (20.03 mg/L) and the wet season's July (12.0 mg/L), when they reach their highest levels. A range of values between 1.6 mg/L and 20.03 mg/L was discovered. Therefore, potassium levels may be caused by farmers using potassium fertilizers, which later settle below and permeate into groundwater. Additionally, potassium might be a naturally occurring resource in the study location.

Additionally, it might be concluded that greater potassium levels during wet seasons may be caused by soil potassium leaking from potassium fertilizers into nearby surface waterways. Meanwhile, the natural occurrence of potassium in rock resulting from some mining-related potassium accounts for the rise in potassium concentration as one approaches the area. Due to the abundance of industrial activity related to a gold mine in the area, this becomes much more significant when we are there. All industrial gold mining facilities must treat their effluent before discharging it into the environment to protect the ecosystem by minimizing the negative effects.




**Figure 28** Monthly variation of K<sup>+</sup> in surface water.

### **3.3 Monthly Variations in Heavy Metal Concentration in Surface Water**

Heavy metal pollution has developed due to human activity, which is the main cause of pollution. This activity frequently results in metal mining, smelting, foundries, and other metals on metal, as well as the leaching of metals from special repositories like landfills, waste dumps, excretion, cattle manure, runoffs, vehicles, and roadwork. The secondary source of heavy metal pollution in the agriculture sector includes the use of herbicides, insecticides, fertilizers, and other heavy metal-containing products. Natural factors such as volcanic activity, metallic corrosion, metallic evaporation from soil and water and sediment re-suspension, soil erosion, and geological weathering can also grow heavy metallic pollutants. In surface waterways, substantial sources of contamination are therefore thought to be trace metals. Trace metal contamination in the surface water is a serious global problem due to its toxicity, ubiquity, and environmental durability.


Seasonal analyses of the amounts and distribution of metals revealed that lead (Pb) concentrations in surface water are shown in Figure 29 for the analyzed surface waters. Pb levels in surface water reached their highest point in December (2.23 mg/L), during the dry season. Nevertheless, the lowest amount in surface water was recorded in February (0.75 mg/L). Thus, the Pb levels are extremely high and would be acutely toxic to virtually all aquatic organisms exposed to this water. Sources of lead contamination include mining, paint, battery waste, coal burning, pesticides, herbicides, and emissions from the burning of leaded fuel. According to research on the health consequences of Pb, metal accumulation in humans would have negative effects on the heart, blood pressure, incidence of hypertension, kidney function, and reproductive issues [1-3]. Children are most susceptible to the harmful effects of Pb, which is a severe hazard to public health. Children's nervous systems and brain development are impacted by Pb hazardous drinking water [12, 78, 79]. According to Brown & Woolf's [80] survey findings in Zamfara, children who live near Pb mines are more likely to develop hemorrhagic encephalopathy due to high Pb levels.

Additionally, Gyamfi *et al.* [81] revelation of increased Pb concentrations in Ghanaian mining site soil and water supported the findings of the current investigation. Therefore, Pb in surface water may come from gold mining, the plastic and rubber, paint, metal, and alloy industries, and battery, fabric, and solid waste disposal industries, among others can reach toxic levels for aquatic organisms. Surface waters nearby get untreated industrial wastewater and sewage from the city. While research on lead exposure in drinking water has been widely documented over the past few decades as lead contamination cases has increased [25], lead continues to be a deadly heavy metal. Nearly every physiological system may be distressed, although the hematologic, gastrointestinal, and neurological systems are most commonly impacted. Furthermore, exposure to lead harms children's behavioral and mental health, making them more susceptible to medical diseases [2, 10, 12, 29, 31, 32].




**Figure 29** Monthly variation of lead (Pb) concentration in surface water.

Figure 30 shows the monthly fluctuations in surface water cadmium (Cd) concentration. Surface water Cd concentrations were generally higher, peaking at 2.73 mg/L in December and declining to 0.061 mg/L in May. Thus, they would be acutely toxic to virtually exposed aquatic organisms. While preparing Cd-Ni batteries, electroplating, control rods, and shields inside nuclear reactors leak Cd into the river, so do stainless steel production facilities, electroplating factories, and vehicle batteries. Under adequate Physicochemical conditions, the predominance of an exchangeable fraction of Cd indicates the anthropogenic origin and strong mobility between aqueous (water) and solid (sediment) phases. Therefore, under proper Physicochemical conditions, the prevalence of an exchangeable fraction of Cd indicates the anthropogenic origin and high mobility between aqueous (water) and solid (sediment) phases [9, 11, 31, 32]. Cadmium is another dangerous element that has been designated as Group B1 by the US EPA (probable human carcinogen). Industrial waste and agricultural fertilizers generate cadmium pollution in drinking water. Renal failure, liver damage, muscle cramps, diarrhea, nausea, and vomiting are a few examples of medical conditions that may be brought on by cadmium exposure [1-10].




**Figure 30** Monthly variation of cadmium (Cd) concentration in surface water.

Figure 31 shows the levels of chromium (Cr) in surface water. Particularly during the dry season and the beginning of the rainy season, the levels of Cr in surface water were greater (May). Surface water chromium concentrations peaked in December (2.09 mg/L) and ranged from 0.32 to 2.09 mg/L. Surface water samples with greater Cr concentrations result from the metal building up over time in the area. According to Fagbenro *et al.* [82], Osun State had greater chromium (Cr) concentration. However, despite exceeding the WHO-recommended level, the concentration of Cr found in this study was not higher than that reported by Fagbenro *et al.* [82].



**Figure 31** Monthly variation of chromium (Cr) concentration in surface water.

Figure 32 shows the manganese (Mn) concentration in surface water. Over the study's twelve months, there was a noticeable variation in focus. December saw the highest Mn levels (1,802) while March saw the lowest (0.35 mg/L). Compared to the rainy season, the concentration of Mn was greater in the dry season. Although manganese is a vital element and is more abundant in surface water, an excessive amount can be detrimental. Our study's high Mn concentration was consistent with Omotola *et al.* [83] reports regarding a gold mining site in Zamfara and Fagbenro *et al.* [82] regarding the heavy metal profile of sediments in gold mining towns in Osun State. Also. Compared to the WHO-recommended standard, Cr and Cd are greater in all samples taken from the study site [68, 69].




**Figure 32** Monthly variation of manganese (Mn) concentration in surface water.

Figure 33 displays the concentrations of mercury (Hg) in surface water. In February (1.39 mg/L), April (1.61 mg/L), August (1.73 mg/L), and December (1.97 mg/L), the concentration of Hg were high. The highest level was recorded in December (1.97 mg/L), while the lowest level (0.13 mg/L)


was recorded in October. Surface water collected along the river's course had an extremely high hg content. This might be due to a long-term buildup of mercury in the sediment of the local river and its tributaries. According to Veiga *et al.* [84], gold mining causes 20–30% of Hg pollution. This is because mercury is used in the process of extracting gold. Water and soil pollution result from the indiscriminate release of metal into water bodies during mining. A higher concentration of Hg was found in the soil taken from various artisanal gold mining sites in Ghana, according to Mantey *et al.* [85]. The findings of their investigation were consistent with those of the current study.

Additionally, the considerable positive association between mercury and temperature supports the finding that mercury toxicity increases as temperature rises [84, 85], leading us to record extraordinarily high mercury levels in water samples. As a result, the discovery of mercury in the surface water suggested that untreated sewers transporting the trash from urban and industrial effluent may have been the cause of the pollution. Therefore, it is crucial to analyze the amounts of mercury in surface water, such as rivers and lakes, because it is a hazardous element with no biological or physiological purpose in humans. Leukemia, neuropathological deterioration, kidney disease, renal system failure, and other health issues are all caused by it, though [29].



**Figure 33** Monthly variation of mercury (Hg) concentration in surface water.

The toxicity of arsenic depends on its chemical state; the inorganic forms are thought to be more dangerous than the organic ones since they have quite different impacts and metabolic processes. Arsenic is a common contaminant throughout the world. Figure 34 shows the amount of arsenic (As) in surface water. Surface water contained As concentrations ranging from 0.04 to 7.36 mg/L. Surface water As levels were consistently low and constant throughout the other months, with a definite high in February (7.37 mg/L). It was interesting to note the elevated As concentrations affected the river and its tributaries. This result was most likely related to anthropogenic activity as well. These heavy metals, however, accumulated throughout the dry season due to human activity in the mining region. Additionally, the discharge of urban and industrial wastewater, particularly sewage from gold mining operations, is to blame for considerable increases in arsenic concentrations in surface water.



**Figure 34** Monthly variation of Arsenic (As) concentration in surface water.

In conclusion, the concentrations of several physicochemical parameters, such as DO, Hardness, Turbidity, Chloride, Potassium, Lead, TSS, Cadmium, Chromium, Manganese, Mercury, and Arsenic, among others, are influenced by the ions (cations and anions). The primary sources of trace elements are mining activities, the aerospace sector, solid rocket fuels, end-of-life vehicle waste, different dyes, and pigments [9-12]. Increased levels of these hazardous metals in the environment reduce agricultural output and soil microbial activity, endangering human health through the food chain. Additionally, these metals may cause problems for human reproduction, biotransformation, and growth [28-32]. Heavy metals accumulate in various plant tissues by interfering with numerous metabolic processes, including inhibition of photosynthesis and respiration and degeneration of main cell organelles. They hurt their growth and development. These effects include stunted growth, delayed germination, chlorosis, premature leaf fall, senescence, decreased crop yield, and loss of enzyme activities [78-80, 86, 87]. Consuming heavy metals like Cd and Zn in humans can lead to various illnesses, including acute gastrointestinal, musculoskeletal, and respiratory problems, as well as harm to the brain, heart, and kidneys [1-10]. Chronic bronchitis, lung cancer, immunotoxicity, neurotoxicity, genotoxicity, infertility, and skin conditions are only a few of the harmful impacts that Ni can produce [11-25]. While excessive Al is extremely neurotoxic for animals and is suspected to be linked to several skeletal abnormalities and neurodegenerative diseases, the toxic effects of Cd include kidney and lung damage, fragile bones, gastrointestinal disorders, carcinogenic, mutagenic, and Itai-Itai disease [9-12].

#### 4. Conclusion and Implications of the Study

The buildup of heavy metals poses long-term repercussions of the threat created by these metals on aquatic life and those who depend on rivers for domestic use and as a source of drinking water. Heavy metals can exist naturally in the environment or through man-made causes. Thus, many governments must set goals to fulfill their pledges to protect water quality to reduce anthropogenic activities by the 2015 Paris Agreement. At the same time, the research region is marked by practices promoting heavy metals buildup. Different heavy metals were found in the surface water samples taken from the study sites, according to the report. The deposition of runoff and garbage from mining sites, residences, agro-processing facilities, and herbicide and pesticide residues from farmers is to blame for heavy metals buildup [33-39]. According to Salazar-

Camacho *et al.* [85], residential, industrial, and mining waste are the main causes of heavy metal pollution in water. High concentrations of elements including Cd, Cr, Hg, As, Pb, Fe, Mn, Ni, Zn, and Cu result from mining. The majority of these substances could be harmful to human health as well as aquatic life. The investigation found dangerous levels of sulfate (SO<sub>2</sub>), nitrate (NO<sub>2</sub>), lead (Pb), cadmium (Cd), chromium (Cr), manganese (Mn), mercury (Hg), and arsenic (As), all of which are determined at high concentrations that are above the WHO limit for drinking water. In Osun State's gold mining towns, soil samples contained several heavy metals in concentrations above WHO-recommended limits. As a result, it has been determined that most surface water used in mining environments is unfit for human consumption. According to the most deviating biological and radiological indicators compared to norms set by reputable local and international organizations, this was attributable to the high amount of contamination from various sources. The relatively high concentration of heavy metal loading in the water is a substantial barrier to rural residents' access to potable water in the mining environment. As a result, the appropriate constituted authority shall continuously monitor the surface water to identify any changes in the water quality. More research on report quality needs to be done in Nigeria's other mining environments to develop technical capacity. In the context of new contaminants and a changing climate, this study suggests additional research directions to enhance knowledge of surface water in a mining setting and sustainable surface water management in mining areas.

## 5. Recommendation

The various findings on the surface water quality in the gold mining areas of Osun State, South-West Nigeria, have been given and addressed in this paper. Therefore, future study should focus on the following issues:

- i. To protect the citizens of this community from the risks of chemical exposure and poisoning brought on by the use of surface or groundwater for irrigation, local authorities like the local government must first offer an alternate supply of water.
- ii. Significant remediation activities are required to clean up the environment, including the soil and water, and artisanal and small-scale miners must be educated on safer and more environmentally friendly techniques of mining. As a result, the environmental concerns connected with the usage and release of hazardous substances like mercury will be reduced.
- iii. Surface water samples from urban, rural, industrial, and remote locations need to have their chemical composition, speciation, and abundance of metal contaminants, as well as their physico-chemical properties, better characterized. To do this, advancements in analytical techniques and equipment are needed.
- iv. In order to improve our understanding of how physico-chemical and heavy metal properties affect surface water processes, we urge further research in the following areas: (a) more characterizations of experiments are needed to provide specifics about the effects of physicochemical and heavy metal properties on surface water; and (b) more research is encouraged because knowledge of these parameters is crucial for more accurate modeling studies and predictions.
- v. Encourage the development of technology for safe and effective water usage, particularly with regard to the reuse and recycling of waste water.

- vi. Encourage private sector investment and the development of suitable water and sanitation technologies as well as waste management infrastructure by promoting environmental health impact analysis (EHIA) as a component of environmental impact assessments (EIA) for all developmental undertakings.

## **Acknowledgements**

The authors are thankful to the Department of Environmental Health Sciences, Kwara State University for the encouragement of research activity and for providing necessary laboratory facilities.

## **Author Contributions**

All of the authors have the same contribution, having read and approved the final manuscript.

## **Funding**

The present research did not receive any financial support.

## **Conflict of Interest**

The authors declare that there is not any conflict of interests regarding the publication of this manuscript. In addition, the ethical issues, including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancy has been completely observed by the authors.

## **References**

1. Raimi M, Ezekwe C. Assessment of trace elements in surface and ground water quality. Mauritius: LAP Lambert Academic Publishing; 2017.
2. Raimi MO, Sawyerr OH, Ezekwe CI, Salako G. Toxicants in water: Hydrochemical appraisal of toxic metals concentration and seasonal variation in drinking water quality in oil and gas field area of rivers state, Nigeria. In: Heavy metals-new insights. London: IntechOpen; 2022.
3. Raimi OM, Ezekwe CI, Bowale A. Statistical and multivariate techniques to trace the sources of ground water contaminants and affecting factors of groundwater pollution in an oil and gas producing wetland in rivers state, Nigeria. medRxiv. 2021. Doi: 10.1101/2021.12.26.21268415.
4. Morufu OR, Henry OS, Clinton IE, Gabriel S. Many oil wells, one evil: Potentially toxic metals concentration, seasonal variation and human health risk assessment in drinking water quality in Ebocha-Obrikom oil and gas area of rivers state, Nigeria. medRxiv 2021. Doi: 10.1101/2021.11.06.21266005.
5. Raimi MO, Sawyerr OH, Ezekwe CI, Olaniyi OA. Quality water not everywhere: Exploratory analysis of water quality across Ebocha-Obrikom oil and gas flaring area in the Core Niger Delta Region of Nigeria [Preprint]. Durham: Research Square LLC.; 2021. Available from: <https://doi.org/10.21203/rs.3.rs-953146/v1>.

6. Afolabi AS, Morufu OR. Investigating source identification and quality of drinking water in Piwoyi Community of Federal Capital Territory, Abuja Nigeria [Preprint]. Durham: Research Square LLC.; 2021. doi: 10.21203/rs.3.rs-736140/v1.
7. Ezekwe CI, Otiasah CL, Raimi MO, Iyingiala AA. Hydrocarbon-based contaminants in drinking water sources and shellfish in the Soku oil and gas fields of South-South Nigeria. *Open J Yangtze Gas and Oil*. 2022; 7: 213-230.
8. Raimi MO, Albert O, Iyingiala AA, Deinkuro NS, Telu M. An environmental scientific report into the crude oil spillage incidence in Tein community, Biseni, Bayelsa State Nigeria. *J Environ Chem Toxicol*. 2022; 6: 01-06.
9. Raimi MO, Ezekwe CI, Bowale A, Samson TK. Hydrogeochemical and multivariate statistical techniques to trace the sources of ground water contaminants and affecting factors of groundwater pollution in an oil and gas producing wetland in rivers state, Nigeria. *Open J Yangtze Oil Gas*. 2022; 7: 166-202.
10. Raimi MO, Sawyerr HO, Ezekwe CI, Opasola AO. Quality water, not everywhere: Assessing the hydrogeochemistry of water quality across Ebocha-Obrikom oil and gas flaring area in the Core Niger Delta Region of Nigeria. *Pollution*. 2022; 8: 751-778.
11. Raimi MO. Preliminary study of groundwater quality using hierarchical classification approaches for contaminated sites in indigenous communities associated with crude oil exploration facilities in rivers state, Nigeria. *Open J Yangtze Oil Gas*. 2022; 7: 124-148.
12. Raimi MO, Sawyerr OH, Ezekwe CI, Salako G. Many oil wells, one evil: Comprehensive assessment of toxic metals concentration, seasonal variation and human health risk in drinking water quality in areas surrounding crude oil exploration facilities in rivers state, Nigeria. *Int J*. 2022; 6: 23-42.
13. Afolabi AS, Raimi MO. When water turns deadly: Investigating source identification and quality of drinking water in Piwoyi Community of Federal Capital Territory, Abuja Nigeria. *Online J Chem*. 2021; 1: 38-58.
14. Raimi MO, Ayibatonbira AA, Anu B, Odipe OE, Deinkuro NS. 'Digging Deeper' evidence on water crisis and its solution in Nigeria for Bayelsa State: A study of current scenario. *Int J Hydro*. 2019; 3: 244-257.
15. Raimi MO, Funmilayo AA, Major I, Odipe OE, Muhammadu IH, Chinwendu O. The sources of water supply, sanitation facilities and hygiene practices in an island community: Amassoma, Bayelsa State, Nigeria. *Public Health Open Access*. 2019; 3: 000134.
16. Raimi MO, Vivien OT, Odipe OE, Owobi OE. The sources of water supply, sanitation facilities and hygiene practices in oil producing communities in central senatorial district of Bayelsa state, Nigeria. *MOJ Public Health*. 2018; 7: 337-345.
17. Raimi MO, Nimsingha D, Odipe OE, Olalekan AS. Health risk assessment on heavy metals ingestion through groundwater drinking pathway for residents in an oil and gas producing area of Rivers State, Nigeria. *Open J Yangtze Oil Gas*. 2018; 3: 191-206.
18. Raimi MO, Ezugwu SC. An Assessment of trace elements in surface and ground water quality in the Ebocha-Obrikom oil and gas producing area of rivers state, Nigeria. *Int J Sci Eng Res*. 2017; 8. Available from: <https://ssrn.com/abstract=3155472>.
19. Raimi MO, Pigha TK, Owobi OE. Water-related problems and health conditions in the oil producing communities in central senatorial district of Bayelsa State. *Imp J Interdiscip Res*. 2017; 3. Available from: <https://ssrn.com/abstract=3108032>.

20. Raimi MO, Ezekwe CI, Sawyerr HO. Problematic groundwater contaminants: Impact of surface and ground water quality on the environment in Ebocha-Obrikom oil and gas producing area of rivers state, Nigeria. 2021. Available from: <https://ssrn.com/abstract=3925848>.
21. Raimi MO. 21st century emerging issues in pollution control. Proceedings of the 6th Global Summit and Expo on Pollution Control; 2019 May 6-7; Amsterdam, Netherlands.
22. Odipe OE, Raimi MO, Suleiman F. Assessment of heavy metals in effluent water discharges from textile industry and river water at close proximity: A comparison of two textile industries from Funtua and Zaria, North Western Nigeria. *Madridge J Agric Environ Sci*. 2018; 1: 1-6.
23. Sawyerr HO, Oluwaseun OE, Asabi OS, Olalekan RM. Bacteriological assessment of selected hand dug wells in students' residential area: A case study of osun state college of health technology, Ilesa, Nigeria. *Glob Sci J*. 2019; 7: 1025-1030.
24. Raimi MO, Adedotun AT, Emmanuel OO, Anu B. An analysis of bayelsa state water challenges on the rise and its possible solutions. *Acta Sci Agric*. 2019; 3: 110-125.
25. Raimi MO, Odipe OE, Anu B, Omini DE, Akpojubaro EH, Owobi OE. Leaving no one behind? Drinking water challenge on the rise in Niger Delta Region of Nigeria: A review. *Merit Res J Environ Sci Toxicol*. 2020; 6: 31-49.
26. Gift RA, Raimi MO, Owobi OE, Oluwakemi RM, Anu B, Funmilayo AA. Nigerians crying for availability of electricity and water: A key driver to life coping measures for deepening stay at home inclusion to slow Covid-19 spread. *Open Access J Sci*. 2020; 4: 69-80.
27. Raimi AA, Raimi MO. Access to electricity and water in Nigeria: A panacea to slow the spread of Covid-19. *Open Access J Sci*. 2020; 4: 34.
28. Raimi MO, Saliu AO, Babatunde A, Okon OG, Taiwo PA, Ahmed AK, et al. The challenges and conservation strategies of biodiversity: The role of government and non-governmental organization for action and results on the ground. In: *Biodiversity in Africa: Potentials, threats and conservation*. Singapore: Springer; 2022. pp. 473-504.
29. Raimi MO, Iyingiala AA, Sawyerr OH, Saliu AO, Ebuete AW, Emberru RE, et al. Leaving no one behind: Impact of soil pollution on biodiversity in the global south: A global call for action. In: *Biodiversity in Africa: Potentials, threats and conservation*. Singapore: Springer; 2022. pp. 205-237.
30. Raimi MO, Abiola I, Alima O, Omini DE. Exploring how human activities disturb the balance of biogeochemical cycles: Evidence from the carbon, nitrogen and hydrologic cycles. *Res World Agric Econ*. 2021; 2: 426.
31. Raimi MO. Assessing pesticides residue in water and fish and its health implications in the Ivo river basin of South-eastern Nigeria. *MOJ Public Health*. 2022; 11: 136-142.
32. Clinton-Ezekwe IC, Osu IC, Ezekwe IC, Raimi MO. Slow death from pollution: Potential health hazards from air quality in the Mgbede oil fields of south-south Nigeria. *Open Access J Sci*. 2022; 5: 61-69.
33. Isah HM, Raimi MO, Sawyerr HO. Probabilistic assessment of self-reported symptoms on farmers health: A case study in kano state for Kura Local Government area of Nigeria. *Environ Anal Ecol Stud*. 2021; 9: 975-985.
34. Hussain MI, Morufu OR, Henry OS. Patterns of chemical pesticide use and determinants of self-reported symptoms on farmers health: A case study in Kano State for Kura local government area of Nigeria. *Res World Agric Econ*. 2021; 2: 342.

35. Morufu OR, Tonye VO, Ogah A, Henry AE, Abinotami WE. Articulating the effect of pesticides use and Sustainable Development Goals (SDGs): The science of improving lives through decision impacts. *Res World Agric Econ*. 2021; 2: 347.
36. Isah HM, Sawyerr HO, Raimi MO, Bashir BG, Haladu S, Odipe OE. Assessment of commonly used pesticides and frequency of self-reported symptoms on farmers health in Kura, Kano State, Nigeria. *J Educ Learn Manage*. 2020; 1: 31-54.
37. Raimi MO, Muhammadu IH, Udensi LO, Akpojubaro EH. Assessment of safety practices and farmers behaviors adopted when handling pesticides in rural Kano state, Nigeria. *Arts Humanit Open Access J*. 2020; 4: 191-201.
38. Muhammadu IH, Raimi MO, Sawyerr OH, Emmanuel OO, Bashir BG, Suleiman H. Qualitative adverse health experience associated with pesticides usage among farmers from Kura, Kano State, Nigeria. *Merit Res J Med Med*. 2020; 8: 432-447.
39. Suleiman RM, Raimi MO, Sawyerr OH. A deep dive into the review of national environmental standards and regulations enforcement agency (NESREA) act. *Int Res J Appl Sci*. 2019; 1: 108-125.
40. Akpanowo MA, Bello NA, Umaru I, Iyakwari S, Joshua E, Yusuf S, et al. Assessment of radioactivity and heavy metals in water sources from Artisanal mining areas of Anka, Northwest Nigeria. *Sci Afr*. 2021; 12: e00761.
41. Corredor JA, González GL, Granados MV, Gutiérrez L, Pérez EH. Use of the gray water footprint as an indicator of contamination caused by artisanal mining in Colombia. *Resour Policy*. 2021; 73: 102197.
42. Masocha M, Dube T, Mambwe M, Mushore TD. Predicting pollutant concentrations in rivers exposed to alluvial gold mining in Mazowe Catchment, Zimbabwe. *Phys Chem Earth*. 2019; 112: 210-215.
43. Ofosu G, Sarpong D. Beyond the doom: Sustainable water management practices of small-scale mining operations. *Resour Policy*. 2022; 77: 102649.
44. Bansah KJ. From diurnal to nocturnal: Surviving in a chaotic artisanal and small-scale mining sector. *Resour Policy*. 2019; 64: 101475.
45. Baffour-Kyei V, Mensah A, Owusu V, Horlu GS. Artisanal small-scale mining and livelihood assets in rural southern Ghana. *Resour Policy*. 2021; 71: 101988.
46. Mensah L. Legal pluralism in practice: Critical reflections on the formalisation of artisanal and small-scale mining (ASM) and customary land tenure in Ghana. *Extr Ind Soc*. 2021; 8: 100973.
47. Ofori AD, Mdee A, Van Alstine J. Politics on display: The realities of artisanal mining formalisation in Ghana. *Extr Ind Soc*. 2021; 8: 101014.
48. Weng L, Margules C. Challenges with formalizing artisanal and small-scale mining in Cameroon: Understanding the role of Chinese actors. *Extr Ind Soc*. 2022; 9: 101046.
49. Kazapoe RW, Amuah EE, Dankwa P. Sources and pollution assessment of trace elements in soils of some selected mining areas of southwestern Ghana. *Environ Technol Innov*. 2022; 26: 102329.
50. Sanchez DN, Knapp CW, Olalekan RM, Nanalok NH. Oil spills in the Niger Delta Region, Nigeria: Environmental fate of toxic volatile organics [Preprint]. Durham: Research Square LLC.; 2021. Available from: <https://doi.org/10.21203/rs.3.rs-654453/v1>.
51. Deinkuro NS, Knapp CW, Raimi MO, Nimlang NH. Environmental fate of toxic volatile organics from oil spills in the Niger Delta Region, Nigeria. *Int J Environ Eng Educ*. 2021; 3: 89-101.

52. Okoyen E, Raimi MO, Oluwatoyin OA, Williams EA. Governing the environmental impact of dredging: Consequences for marine biodiversity in the Niger Delta Region of Nigeria. *Insights Min Sci Technol.* 2020; 2: 555586.
53. Olalekan RM, Omidiji AO, Williams EA, Christianah MB, Modupe O. The roles of all tiers of government and development partners in environmental conservation of natural resource: A case study in Nigeria. *MOJ Ecol Environ Sci.* 2019; 4: 114-121.
54. Worlanyo AS, Jiangfeng L. Evaluating the environmental and economic impact of mining for post-mined land restoration and land-use: A review. *J Environ Manage.* 2021; 279: 111623.
55. Raimi MO, Ezugwu SC. Influence of organic amendment on microbial activities and growth of pepper cultured on crude oil contaminated Niger delta soil. *Int J Econ Energy Environ.* 2017; 2: 56-76.
56. Sawyerr HO, Raimi MO, Adeolu AT, Odipe OE. Measures of harm from heavy metal pollution in battery technician within Ilorin Metropolis, Kwara State, Nigeria. *Commun Soc Media.* 2019; 2: 73-89.
57. Obiri-Yeboah A, Nyantakyi EK, Mohammed AR, Yeboah SI, Domfeh MK, Abokyi E. Assessing potential health effect of lead and mercury and the impact of illegal mining activities in the Bonsa River, Tarkwa Nsuaem, Ghana. *Sci Afr.* 2021; 13: e00876.
58. Mulenga C. Soil governance and the control of mining pollution in Zambia. *Soil Secur.* 2022; 6: 100039.
59. Søndergaard J, Mosbech A. Mining pollution in Greenland-the lesson learned: A review of 50 years of environmental studies and monitoring. *Sci Total Environ.* 2021; 812: 152373.
60. Adegbola IP, Aborisade BA, Adetutu A. Health risk assessment and heavy metal accumulation in fish species (*Clarias gariepinus* and *Sarotherodon melanotheron*) from industrially polluted Ogun and Eleyele Rivers, Nigeria. *Toxicol Rep.* 2021; 8: 1445-1460.
61. Gbedzi DD, Ofosu EA, Mortey EM, Obiri-Yeboah A, Nyantakyi EK, Siabi EK, et al. Impact of mining on land use land cover change and water quality in the Asutifi North District of Ghana, West Africa. *Environ Challenges.* 2022; 6: 100441.
62. Taiwo AM, Awomeso JA. Assessment of trace metal concentration and health risk of artisanal gold mining activities in Ijeshaland, Osun State Nigeria—Part 1. *J Geochem Explor.* 2017; 177: 1-10.
63. World Health Organization. Progress on sanitation and drinking water: 2015 Update and MDG Assessment [Internet]. WHO/UNICEF Joint Monitoring Programme; 2015. Available from: [https://apps.who.int/iris/bitstream/handle/10665/177752/9789241509145\\_eng.pdf?sequence=1&isAllowed=y](https://apps.who.int/iris/bitstream/handle/10665/177752/9789241509145_eng.pdf?sequence=1&isAllowed=y).
64. Raimi MO, Bilewu OO, Adio Z, Halimat A. Women contributions to sustainable environments in Nigeria. *J Sci Res Allied Sci.* 2019; 5: 35-51.
65. Raimi MO, Suleiman RM, Odipe OE, Tolulope SJ, Modupe O, Olalekan AS, et al. Women role in environmental conservation and development in Nigeria. *Ecol Conserv Sci.* 2019; 1. Doi: 10.2139/ssrn.3425832.
66. Edet AE, Offiong OE. Evaluation of water quality pollution indices for heavy metal contamination monitoring. A study case from Akpabuyo-Odukpani area, Lower Cross River Basin (southeastern Nigeria). *GeoJournal.* 2002; 57: 295-304.
67. APHA. Standard methods for the examination of water and waste water. 17th ed. American Public Health Association. Washington, DC: American Public Health Association; 2012.

68. Herschy RW. Water quality for drinking: WHO guidelines. In: Encyclopedia of earth sciences series. Dordrecht: Springer; 2011. pp. 876-883.
69. World Health Organization. Guidelines for drinking water quality—Fourth edition incorporating the first addendum. Geneva: World Health Organization; 2017.
70. Nwankwo CN, Ogagarue DO. Effects of gas flaring on surface and ground waters in Delta State, Nigeria. *J Geol Min Res*. 2011; 3: 131-136.
71. Egbueri JC, Mgbenu CN, Chukwu CN. Investigating the hydrogeochemical processes and quality of water resources in Ojoto and environs using integrated classical methods. *Model Earth Syst Environ*. 2019; 5: 1443-1461.
72. Raimi MO. Self-reported symptoms on farmers health and commonly used pesticides related to exposure in Kura, Kano State, Nigeria. *Ann Community Med Public Health*. 2021; 1: 1002.
73. Raimi MO. Health risk exposure to cypermethrin: A case study of kano state, Nigeria. *J Agric*. 2018; 1: 14-15.
74. Raimi OM, Samson TK, Sunday AB, Olalekan AZ, Emmanuel OO, Jide OT. Air of uncertainty from pollution profiteers: Status of ambient air quality of sawmill industry in Ilorin Metropolis, Kwara State, Nigeria. *Res J Ecol Environ Sci*. 2021; 1: 17-38.
75. Raimi MO, Adio Z, Emmanuel OO, Samson TK, Ajayi BS, Ogunleye TJ. Impact of sawmill industry on ambient air quality: A case study of Ilorin Metropolis, Kwara State, Nigeria. *Energy Environ Sci*. 2020; 3. Doi: 10.2139/ssrn.3586971.
76. Olalekan RM, Timothy AA, Enabulele Chris E, Olalekan AS. Assessment of air quality indices and its health impacts in Ilorin Metropolis, Kwara State, Nigeria. *Sci Park J Sci Res Impact*. 2018; 4: 060-074.
77. Edna Ateboh P, Raimi MO. Corporate civil liability and compensation regime for environmental pollution in the Niger Delta. *Int J Recent Adv Multidiscip Res*. 2018; 5: 3870-3893.
78. Mohanta VL, Singh S, Mishra BK. Human health risk assessment of fluoride-rich groundwater using fuzzy-analytical process over the conventional technique. *Groundw Sustain Dev*. 2020; 10: 100291.
79. Kochian LV, Piñeros MA, Liu J, Magalhaes JV. Plant adaptation to acid soils: The molecular basis for crop aluminum resistance. *Annu Rev Plant Biol*. 2015; 66: 571-598.
80. Brown MJ, Woolf AD. Chapter 1.8—Zamfara gold mining lead poisoning disaster—Nigeria, Africa, 2010. In: History of modern clinical toxicology. Cambridge: Academic Press; 2022. pp. 97-107.
81. Gyamfi E, Appiah-Adjei EK, Adjei KA. Potential heavy metal pollution of soil and water resources from artisanal mining in Kokoteasua, Ghana. *Groundw Sustain Dev*. 2019; 8: 450-456.
82. Fagbenro AA, Yinusa TS, Ajekiigbe KM, Oke AO, Obiajunwa EI. Assessment of heavy metal pollution in soil samples from a gold mining area in Osun State, Nigeria using proton-induced X-ray emission. *Sci Afr*. 2021; 14: e01047.
83. Oladipo MO, Njinga RL, Elele UU, Salisu A. Heavy metal contaminations of drinking water sources due to illegal gold mining activities in Zamfara state-Nigeria. *J Chem Biochem*. 2014; 2: 31-44.
84. Veiga MM, Maxson PA, Hylander LD. Origin and consumption of mercury in small-scale gold mining. *J Clean Prod*. 2006; 14: 436-447.

85. Mantey J, Nyarko KB, Owusu-Nimo F, Awua KA, Bempah CK, Amankwah RK, et al. Influence of illegal artisanal small-scale gold mining operations (galamsey) on oil and grease (O/G) concentrations in three hotspot assemblies of Western Region, Ghana. *Environ Pollut.* 2020; 263: 114251.
86. Boening DW. Ecological effects, transport, and fate of mercury: A general review. *Chemosphere.* 2000; 40: 1335-1351.
87. Salazar-Camacho C, Salas-Moreno M, Marrugo-Madrid S, Paternina-Uribe R, Marrugo-Negrete J, Díez S. A human health risk assessment of methylmercury, arsenic and metals in a tropical river basin impacted by gold mining in the Colombian Pacific region. *Environ Res.* 2022; 212: 113120.